



Actions required to secure the large-scale deployment of the leading CDR approaches to meet EU climate targets

## C-SINK– FACTSHEET LECTURE 1 DELIVERED BY ICAMCYL & EKOLIVE

### CARBON DIOXIDE REMOVAL TECHNOLOGIES (CDR-T): BIOLOGICAL CO<sub>2</sub> FIXATION (BCO<sub>2</sub>)

#### INTRODUCTION

In the Paris Agreement (United Nations, 2015), the necessary bases were established to address the issue of climate change as urgent, using the highest level of technological and scientific knowledge to contain and address the problem on a global scale. Subsequently, the Intergovernmental Panel on Climate Change (IPCC) announced the results of a rigorous assessment based on scientific, technical, and socio-economic data available in the literature on global warming of 1.5°C and the comparison between global warming of 1.5°C and 2°C above pre-industrial levels. Resulting model pathways with no or limited overshoot of 1.5°C indicate that emissions must decline by 45% to keep warming below 1.5°C by 2030, and towards net zero by 2050 (IPCC, 2018b). The C-SINK project funded by Horizon Europe aims to establish the foundations to build a standardized and transparent European CDR market with trustworthy accounting methodologies based on robust Monitoring, Reporting, and Verification (MRV) pre-standards and policy strategies.

#### WHAT IS CARBON DIOXIDE REMOVAL (CDR)?

Following the IPCC's 8<sup>th</sup> Assessment Report, CDR refers to activities that remove carbon dioxide (CO<sub>2</sub>) from the atmosphere and store it permanently (IPCC, 2018a). They can be grouped into nature-based and technology-based removals (Meyer-Ohlendorf, 2020). Nature-based enhancing biological sinks of CO<sub>2</sub> are, among others, afforestation (AF), reforestation (RF), and soil carbon sequestration (SCS) as a result of BCO<sub>2</sub>. Technology-based removals employ chemical engineering to achieve long-term removal and storage. Among others, an example is bioenergy with carbon capture and storage (BECCS), but there are

additional approaches including improving soil quality, enhanced weathering, and the production of biochar (Ornelas et al., 2023).



#### GLOSSARY

**Aerobic metabolism:** Type of metabolism which occurs in the presence of oxygen.

**Anaerobic metabolism:** Type of transformation produced with no presence of oxygen.

**Autotrophic bacteria:** Type of bacteria which synthesizes their own food. They derive energy from light, utilize simple inorganic compounds like CO<sub>2</sub>, water, etc., and convert them into organic compounds to supplement their energy requirements.

**Biological carbon fixation (BCO<sub>2</sub>):** A nature-based technology in which living organisms absorb and convert CO<sub>2</sub> into organic matter through biochemical pathways (Berg, 2011). Organic compounds created by BCO<sub>2</sub> are used to store energy and to build other biomolecules. Carbon is primarily fixed through photosynthesis, when sunlight is available, and chemosynthesis, in the absence of sunlight.

**Heterotrophic microorganisms:** Organisms that obtain their energy and carbon from organic compounds.

**Soil microbiota:** The abundance and diversity of microbes residing in the soil (bacteria, fungi, archaea, protists, and algae).

#### DEFINING CO<sub>2</sub> REMOVAL FROM A BCO<sub>2</sub> STANDPOINT & PROCESS


CO<sub>2</sub> removal by BCO<sub>2</sub> refers to the amount of CO<sub>2</sub> that is consumed and stored by microorganisms in the soil and plants growing in a defined area. When plants photosynthesize, they take CO<sub>2</sub> from the atmosphere and convert it into organic



Funded by  
the European Union

Funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Climate, Infrastructure and Environment Executive Agency (CINEA). Neither the European Union nor CINEA can be held responsible for them.

C-SINK– FACTSHEET LECTURE 1 DELIVERED BY ICAMCYL & EKOLIVE



matter, such as wood, leaves, and roots, while releasing oxygen (see Figure 1). Apart from CO<sub>2</sub> fixation by plants, BCO<sub>2</sub> occurs in all microorganisms, both autotrophic and heterotrophic. Biological carbon fixation by autotrophs can utilize light or inorganic chemical energy to fix atmospheric CO<sub>2</sub> through evolved carbon fixation pathways (Gong et al., 2018). Autotrophic bacteria synthesize all their constituent cells using CO<sub>2</sub> as a carbon source in that way (Ge et al., 2016). Importantly, recent studies report significant biological CO<sub>2</sub> fixation by heterotrophic microorganisms. These microorganisms are capable of consuming CO<sub>2</sub> and incorporating it into their metabolism during bioremediation and biofertilization processes (Bräuer et al., 2016); (Chen et al., 2009); (Kimura et al., 2011); (Miltner et al., 2005); (Santoro et al., 2013).

**WHAT IS THE CURRENT STATE OF THE TECHNOLOGY?**

Various microorganism based biostimulants are commonly applied by farmers to increase yields – however, impact on CO<sub>2</sub> fixation in soil was never the main purpose of

their use, consequently monitoring and verification of their impact are lacking. The major unknowns in BCO<sub>2</sub> technology are the poorly understood related processes involved, the scalability, and the very limited data. There are no clear MRV standards or methodologies. It is very well known that carbon gets sequestered in soil as Soil Organic Carbon (SOC), but there is also a possibility of carbon sequestration in the form of Soil Inorganic Carbon (SIC). Moreover, there is a lack of understanding of biochemical pathways happening in soil microbiota. Furthermore, methodologies for the determination of carbon sequestered in soil are not uniform and standardized at the moment. SIC is not taken into account when calculating CO<sub>2</sub> stored in the form of soil carbon.

**DEFINITION OF EACH PROCESS-STEP**

1. Biobleaching or microbiological weathering for 5 days, in which bacteria multiply and produce metabolites dissolving minerals, to produce 2 types of biostimulants: ekofertile® plant and microfertile® plant - water consumption.

## C-SINK– FACTSHEET LECTURE 1 DELIVERED BY ICAMCYL & EKOLIVE

2. Pumping of biostimulants to plastic canisters – *energy consumption by the pump.*
3. Printing the labels - *energy consumption.*
4. Transport of biostimulants to farmers – *fuel consumption.*
5. The farmer mixes biostimulants with water and applies mixed biostimulants by spraying – *fuel consumption, water consumption, or by irrigation system – electricity and water consumption.*
6. Organic and inorganic carbon monitoring in soil.
7. Harvest and analysis of yield – harvest by tractors, transport to food factories – *fuel consumption.*




Figure 2. BCO<sub>2</sub> step-process.

### PERMANENCE

The most common timeframe applied in BCO<sub>2</sub> in the form of SCS is decades to centuries according to IPCC1. Regarding the time factor in removing CO<sub>2</sub>, soil microbiota communities show significant roles in carbon removal from the atmosphere and fixing it in the soils. However, potential fluctuations during large-scale deployment regarding microbial communities have not been sufficiently studied. Carboxylase reactions are rather immediate reactions and result in the assimilation of CO<sub>2</sub> into organic metabolites by microorganisms. This CO<sub>2</sub> incorporation by chemo-organo-heterotrophic microorganisms has been referred to as “heterotrophic carbon dioxide assimilation” or “heterotrophic carbon dioxide fixation” and can be assumed to have an immediate effect (Braun et al., 2021).

### REVERSAL RISK

BCO<sub>2</sub> produces SIC and SOC. Regarding SIC, BCO<sub>2</sub> presents a low risk. Sylvera<sup>2</sup> has ranked mineralized CO<sub>2</sub> (SIC) to have no practical risk of reversal. Microbial carbon fixation of SIC is proven in research done by Liu et al. (2020) where they demonstrated that desert soil microbes are directly involved in fixing atmospheric CO<sub>2</sub> via biomineralisation. Same is proven by Braun et al. (2021) for heterotrophic bacteria’s ability to fixate CO<sub>2</sub>. However, Sylvera considers that SOC has a high risk of reversal since there is a high range of factors that trigger carbon reversal — flooding, droughts, wildfires, disease, renovation of a pasture, change in soil acidity or change of the soil nutrient content (named “depletion event” in a guide<sup>3</sup> to the estimating sequestration of carbon in soil using default values method).

### COSTS, TRL, AND TYPICAL SCALE

Some estimated costs are 1,09 € per litre of biostimulant and it is applied based on the type of crops in rates from 10 – 200 litres per hectare. It is affordable for farmers because the cost return of this novel technology is 10 times higher compared to conventional products due to no diseases, yield increase, proteins increase, and possible reduction on N fertilizer up to 50 %. The potential additional profits from carbon credits will be calculated during the field trials within the C-SINK project. Usually, it ranges from an additional 30 – 300 € per hectare.

### TYPICAL SCALE OF CO<sub>2</sub> REMOVAL/LIMITING FACTORS

According to Shimmel (1987), dark fixation of CO<sub>2</sub> in agricultural soil, based on the rate of radiolabelled CO<sub>2</sub> incorporation, varied from 0.2 to 4.8 mg/m<sup>2</sup>h. Data from Šantrůčková et al. (2005) projected the rate of estimated heterotrophic CO<sub>2</sub> fixation which ranged from 36.5 mg CO<sub>2</sub>/m<sup>2</sup>h, in the soil with pH 7.5, to 2.8 mg CO<sub>2</sub>/m<sup>2</sup>h, in the soil with pH 4.8.

<sup>1</sup> [https://www.ipcc.ch/report/ar6/wg3/downloads/outreach/IPCC\\_AR6\\_WGIII\\_Factsheet\\_CDR.pdf](https://www.ipcc.ch/report/ar6/wg3/downloads/outreach/IPCC_AR6_WGIII_Factsheet_CDR.pdf) <sup>2</sup> <https://cer.gov.au/document/guide-to-estimating-sequestration-carbon-soil-using-default-values-method> <sup>3</sup> <https://www.sylvera.com/blog/permanence-carbon-credits>



Actions required to secure the large-scale deployment of the leading CDR approaches to meet EU climate targets

## C-SINK– FACTSHEET LECTURE 1 DELIVERED BY ICAMCYL & EKOLIVE

No industrial-scale data is available. Thereby a pilot is essential to study BCO<sub>2</sub> deployment capacity.

C-SINK experts have found that limiting factors to scaling BCO<sub>2</sub> include obtaining import permissions, local registrations, and sales for the biostimulants in the EU<sup>4</sup>.

### NEGATIVE EFFECTS

High water consumption.

### CO-EFFECTS

Benefits for farmers: sugar and protein content increase, faster development of crops, disease resistance, drought resistance, yield and profit increase with low investment per hectare, increased efficiency of agrochemicals.

Benefit for mining companies: new ecological processing technology increases the value and purity of minerals.

The co-effects described above have been verified and measured<sup>5</sup>, for instance, if the right dosage of the biostimulant is used. Also, the mentioned co-effects can potentially positively influence the use of biostimulants as BCO<sub>2</sub> CDR-T because farmers are primarily interested in applying it due to agricultural benefits and not specifically CO<sub>2</sub> sequestration potential.

The C-SINK project will allow demonstration of biostimulants' role in BCO<sub>2</sub> CDR-T which will further raise awareness and willingness of, not only farmers, but all land managers to apply this technology. This will enable carbon sequestration and contribute to the goals of the Paris Agreement in tackling climate change by supplementing existing CO<sub>2</sub> mitigation efforts.

### BCO<sub>2</sub> PILOTS DEVELOPED WITHIN THE C-SINK PROJECT

Several case studies on farmland with and without crops will be conducted in three countries (Germany, Slovakia, and Croatia). The soil will be inoculated with up to 1000 microbial species of heterotrophic, autotrophic microorganisms and microalgae

produced by bioleaching operations. The objective of case studies will be to evaluate the potential and mechanisms by which microorganisms contribute to soil carbon sequestration and retention. Each location is characterised by different climates and different soil types.

Field trials in Slovakia are conducted on raspberries and cereals, on red currants and blueberries in Germany and, in vineyards in Croatia.

Soil samples are taken periodically throughout the growing season to assess the



Figure 3. C-SINK BCO<sub>2</sub> pilots

soil sequestration potential. This is done according to ISO 18400-104:2018 Soil Quality – Sampling. Soil samples' collection and application of different concentrations of ekofertile<sup>®</sup> plant and microfertile<sup>®</sup> plant to raspberries by spraying and irrigation are shown in Figure 3. - results can be seen at the farm in Košické Olšany. Furthermore, soil samples are analysed to obtain

<sup>4</sup> In EU countries – in some cases it is a simple two-page declaration, and labels submitted to a relevant authority. In other countries no need for anything, it is automatically mutually recognized. E.g. in Spain, it is on a volunteer basis to

prepare this declaration. Only in Switzerland registration is needed, while UK is free to sell without any action.

<sup>5</sup> It is possible to measure yield increase, protein increase etc. to calculate an increase.



## C-SINK– FACTSHEET LECTURE 1 DELIVERED BY ICAMCYL & EKOLIVE

parameters of interest: total organic carbon, total inorganic carbon, humus, pH, total nitrogen, phosphorous, potassium, magnesium, copper, C/N ratio, and bulk density.

To process data and determine SCS, a variety of methodologies will be used to evaluate their correctness and suitability such as Microbial Carbon Mineralization (MCM) - Methodology for Quantification and Crediting of Carbon Dioxide Removal from Andes, VM0042 – Methodology for Improved Agricultural Land Management from Verra, Carbon Credits (Carbon Farming Initiative— Estimation of Soil Organic Carbon Sequestration using Measurement & Models) Methodology Determination 2021 from Australian Government & Soil Carbon Measurement Methodology from Carboneg. Finally, the impact on biological carbon sequestration in the soil will be monitored and compared in sprayed and irrigated areas within the C-SINK project.

### REFERENCES

❖ Berg, I. A. (2011). Ecological aspects of the distribution of different autotrophic CO<sub>2</sub> fixation pathways. *Applied and Environmental Microbiology*, 77(6), 1925–1936. <https://doi.org/10.1128/AEM.02473-10>

❖ Bräuer, S. L., Kranzler, K., Goodson, N., Murphy, D., Simon, H. M., Baptista, A. M., & Tebo, B. M. (2016). Dark Carbon Fixation in the Columbia River's Estuarine Turbidity Maxima: Molecular Characterization of Red-Type cbbL Genes and Measurement of DIC Uptake Rates in Response to Added Electron Donors. *Estuaries and Coasts*, 1–23. <https://doi.org/10.1007/s12237-013-9603-6>

❖ Braun, A., Spona-Friedl, M., Avramov, M., Elsner, M., Baltar, F., Reinthaler, T., Hemdl, G. J., & Griebler, C. (2021). Reviews and syntheses: Heterotrophic fixation of inorganic carbon - Significant but invisible flux in environmental carbon cycling. *Biogeosciences*, 18(12), 3689–3700. <https://doi.org/10.5194/bg-18-3689-2021>

❖ Chen, Y., Wu, L., Boden, R., Hillebrand, A., Kumaresan, D., Moussard, H., Baciu, M., Lu, Y., Colin Murrell, J., & T1. (2009). Life without light: microbial diversity and evidence of sulfur- and ammonium-based chemolithotrophy in Movile Cave. *JOUR AU*, 3(9). <https://doi.org/10.1038/ismej.2009.57>

❖ Ge, T., Wu, X., Liu, Q., Zhu, Z., Yuan, H., Wang, W., Whiteley, A. S., & Wu, J. (2016). Effect of simulated tillage on microbial autotrophic CO<sub>2</sub> fixation in paddy and upland soils. *Scientific Reports*, 6(December 2015), 1–9. <https://doi.org/10.1038/srep19784>

❖ Gong, F., Zhu, H., Zhang, Y., & Li, Y. (2018). Biological carbon fixation: From natural to synthetic. *Journal of CO<sub>2</sub> Utilization*, 28, 221–227. <https://doi.org/10.1016>

❖ IPCC. (2018a). Annex I: Glossary [Matthews, J.B.R. (ed.)]. In: *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the . In Global Warming of 1.5°C.* (pp. 541–562). Cambridge University Press. <https://doi.org/10.1017/9781009157940.008>

❖ IPCC. (2018b). Summary for Policymakers. In: *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to. In Global Warming of 1.5°C.* Cambridge University Press. <https://doi.org/10.1017/9781009157940.001>

❖ Kimura, S., Bryan, C. G., Hallberg, K. B., & Johnson, D. B. (2011). Biodiversity and geochemistry of an extremely acidic, low-temperature subterranean environment sustained by chemolithotrophy. *Environmental Microbiology*, 13(8). <https://doi.org/10.1111/j.1462-2920.2011.02434.x>

❖ Liu, Z., Sun, Y., Zhang, Y., Qin, S., Sun, Y., Mao, H., & Miao, L. (2020). Desert soil sequesters atmospheric CO<sub>2</sub> by microbial mineral formation. *Geoderma*, 361, 114104. <https://doi.org/10.1016/j.geoderma.2019.114104>

❖ Meyer-Ohendorf, N. (2020). *EU Framework for CO<sub>2</sub> Removals-Targets and Commitments.*

❖ Miltner, A., Kopinke, F.-D., Kindler, R., Selesi, D., Hartmann, A., & Kästner, M. (2005). Non-photosynthetic CO<sub>2</sub> fixation by soil microorganisms. *JOUR*, 269(1), 1–23. <https://doi.org/10.1007/s11104-004-0483-1>

❖ Ornelas, A., Mumbai, Y. B., Gautier, P., Lai, P., Paris, L., & London, S. R. (2023). *Carbon Capture, Removal, And Credits Pose Challenges For Companies.* [spglobal.com/ratings](https://spglobal.com/ratings)

❖ Santoro, A. L., Bastviken, D., Gudas, C., Tranvik, L., & Enrich-Prast, A. (2013). Dark Carbon Fixation: An Important Process in Lake Sediments. *PLoS ONE*, 8(6), 1–7. <https://doi.org/10.1371/journal.pone.0065813>

❖ Šantrůčková, H., Bird, M. I., Elhottová, D., Novák, J., Pícek, T., Šimek, M., & Tykva, R. (2005). Heterotrophic Fixation of CO<sub>2</sub> in Soil. *Microbial Ecology*, 49(2). <https://doi.org/10.1007/s00248-004-0164-x>

❖ Shimmel, S. (1987). Dark fixation of carbon dioxide in an agricultural soil. *Soil Science*, 144, 20–23. [https://journals.lww.com/soilsci/fulltext/1987/07000/dark\\_fixation\\_of\\_carbon\\_dioxide\\_in\\_an\\_agricultural.4.aspx](https://journals.lww.com/soilsci/fulltext/1987/07000/dark_fixation_of_carbon_dioxide_in_an_agricultural.4.aspx)

❖ Tavi, D. (2023). *Ekofertile and microfertile plant biostimulants potentials on soil quality and wheat nutrition enhancement.*

❖ United Nations. (2015). *ADOPTION OF THE PARIS AGREEMENT - Paris Agreement text English.* [https://unfccc.int/sites/default/files/english\\_paris\\_agreement.pdf](https://unfccc.int/sites/default/files/english_paris_agreement.pdf)

👉 If you want to know more about the C-SINK project, visit our website at <https://c-sinkproject.eu/>